

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application. Follow all applicable directions, restrictions and precautions on the main EPA registered label and this supplemental label.

This product can be used for forestry sites, including conifer site preparation areas. Not for use in recreational sites, Christmas tree plantations or in ornamental plantings.

GENERAL INFORMATION:

Mountain beavers (*Aplodontia rufa*) are active for short durations throughout the day, but are usually more active in the evening. Surface activity is usually fairly close to their burrows, although they can range over several acres if food is not plentiful. Mountain beaver burrows can be located by observing feeder holes or freshly pushed dirt. There may be several entrances to the mountain beaver burrow system. Mountain beavers exhibit caching and hoarding behavior and are likely to move the bait packet into their burrow. Abandoned burrows are frequently reinvaded by mountain beavers from surrounding areas, and cached bait may control invading animals. Population monitoring, trapping and baiting are components of an Integrated Pest Management (IPM) approach to mountain beaver control. Research has found that trapping in conjunction with baiting is the most effective method of control.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, birds, fish and aquatic invertebrates. Predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have been poisoned by the product. Collect and properly dispose of dead animals. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of empty container in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO NONTARGET ORGANISMS**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**FIFRA Special Local Need
Supplemental Label**

This label must be in the possession of the user at the time of pesticide application. Follow all other applicable directions, restrictions and precautions on the EPA registered label.

**FOR DISTRIBUTION AND USE ONLY IN WITHIN
THE STATE OF OREGON**

(West of the Crest of the Cascade Mountain Range)

This label valid until December 31, 2011 or until otherwise amended, withdrawn, canceled or suspended.

**rozol®
Pellets**

**FOR CONTROL OF MOUNTAIN
BEAVER IN FORESTRY
PLANTATIONS**

Active ingredient: Chlorophacinone0.005%
Inert ingredients:99.995%
Total100.000%

EPA Reg. No. 7173-151
EPA Est. No. 7173-WI-01
EPA SLN No. OR-060026
Label No. 155-0001-1006

KEEP OUT OF REACH OF CHILDREN

CAUTION: See side panel for further cautions and first aid instructions

24(c) Registrant:

LIPHA/TECH®

Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209
(414) 351-1476

Net Weight:

18 lbs (8.8 Kg) (24 x 12 oz. packs)

Use Rate / Bait Placement:

Place one unopened packet of bait inside a feeder hole or runway of an active mountain beaver burrow system. Do not place more than one packet per hole or individual runway; or more than two packets of bait per burrow system. Insert bait packet at least 12 inches into hole or runway. Do not cut or tear open the packets of bait.

Application Timing:

Applications made between October and February have been found to be most effective at reducing damage to tree seedlings planted in late winter or early spring. Applications should be made to target the adult mountain beaver population. Juvenile mountain beavers are more commonly present during the summer months. They are smaller, more naive, and tend to be more active outside of the burrow. To minimize risks to non-target predators and scavengers, do not apply bait from mid-May to September 1.

RESTRICTIONS / PRECAUTIONS:

Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application.

Do not apply to any site, against any pest, or by any method of application not stated on this label.

For use to control mountain beavers only.

Use of Rozol under this Special Local Need (SLN) label is restricted to western Oregon only (West of the crest of the Cascade Mountain Range).

Do not apply this product from mid-May to September 1 when juvenile mountain beavers are present, due to concerns with secondary poisoning of non-target animals.

This pesticide is toxic to fish and wildlife. Rozol Pellets should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may contact the Oregon Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding species listed as threatened or endangered.

Consult the Federal label for additional restrictions and precautions to protect aquatic and other non-target organisms.

Do not apply this product directly to water or within 10 feet of surface water (i.e. streams, rivers, lakes).

Pick up and properly dispose of any spilled bait that cannot be recovered for use.

Keep away from humans, domestic animals and pets during application, handling and storage. (29306)